

# DRUG FACTS

Active ingredient(s)

Purpose

Use(s)  
Uses  
Temporarily relieves minor aches and pains associated with

- # headache
- # backache
- # common cold
- # minor arthritis pain
- # toothache
- # menstrual cramps
- # muscular aches

Temporarily reduces fever.

## Warnings

Do not use  
Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if you have  
Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are  
Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product  
When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if  
Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding

Keep out of reach of children  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

Directions

- do not use more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

## Other information

Drug Facts  
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- rash
- blisters
- skin reddening
- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Adults and children: (12 years and older)  
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:  
Do not give to children under 12 years of age.

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets

- do not use any opened or torn packets

***Inactive ingredients***

***Inactive ingredients***

carnauba wax<sup>\*</sup>, cellulose<sup>\*</sup>, colloidal silicon dioxide, corn starch<sup>\*</sup>, hypromellose, iron oxide red<sup>\*</sup>, lactose, magnesium stearate, microcrystalline cellulose<sup>\*</sup>, polydextrose, polyethylene glycol, povidone, silica<sup>\*</sup>, sodium lauryl sulfate<sup>\*</sup>, sodium starch glycolate, stearic acid<sup>\*</sup>, titanium dioxide, triacetin<sup>\*</sup>  
<sup>\*</sup> may contain